

<h1>25E18</h1> <p>2.5 RM</p>	Trait Options		Geographic Adaptation			PRO-FIT Placement			
		Western West of Miss. River	HR	Moves South well for early season variety	HR	Raising the bar for yield potential!			
		Central (I-35 to I-65)	HR			Resistant to Stem Canker, BSR and Frogeye			
		Eastern East of Miss. River	HR	Moves North well for full season variety	HR	Best of class IDC			
					Can cover lots of acres and environments				
General Characteristics			Agronomic Features			Disease Tolerance			
Plant Height	Med-Tall	Yield for Maturity	Excellent		Brown Stem Rot	Excellent			
Canopy/Plant type	Medium	Emergence	Excellent		Frogeye Leaf Spot	Excellent			
Growth Habit	Medium	Standability	Very Good		Iron Chlorosis	Excellent			
Flower Color	Purple	Shatter Resistance	Very Good		Phytophthora Field Tol.	Very Good			
Hilum Color	Imp.Black	Stress Tolerance	Excellent		SDS Tolerance	Good			
Pod Color	Brown	SCN Resistance Source	PI88788		White Mold	Good			
Pubescence	Gray	Phytophthora Res. Source	None		Stem Canker	Excellent			
Protein	NA	No-Till Adaptability	0		Root Knot	NA			
Oil	NA				Cover Crop Companion	Good			
Soil Placement		Recommended Seeding Rates				Herbicide Tolerance			
Light Soils: Low O.M., Low CEC, Low water holding capacity.	HR	Row Width	120/130	135/145	150/160	Conventional		Dicamba	
Medium Soils: O.M. 1.5-3.5%, CEC 11-20, Good productivity, well drained silt loams.	HR	30" Rows	HR	HR	R	Glyphosate		2,4-D	
Heavy Soils: High O.M. >3.5%, CEC >20, Well drained highly productive with deep top soil	HR	Twin Rows	HR	HR	R				
Poorly Drained Soils: Soils that tend to remain saturated for extended periods of time.	HR	20" Rows	R	R	HR	Glufosinate		Callisto	
		<20" rows	NR	R	HR	Sulfonurea		Isoxoflutol	
<small>Population ranges are listed in 1000 seeds/acre</small>									
NR=Not Recommended, R=Recommended, HR=Highly Recommended									
Notes:									

