

<h1>30E11</h1> <p>3.0 RM</p>	Trait Options		Geographic Adaptation			PRO-FIT Placement			
		Western West of Miss. River	HR	Moves South well for early season variety	HR	High yields, disease tolerance AND standability!			
		Central (I-35 to I-65)	HR			Good on SDS and SWM			
		Eastern East of Miss. River	HR	Moves North well for full season variety	HR	Consistent performer and stands great			
					Works on irrigated acres also				
General Characteristics			Agronomic Features			Disease Tolerance			
Plant Height	Med-Tall	Yield for Maturity	Excellent	Brown Stem Rot	Excellent				
Canopy/Plant type	Med-Bush	Emergence	Excellent	Frogeye Leaf Spot	NA				
Growth Habit	Branchy	Standability	Excellent	Iron Chlorosis	Very Good				
Flower Color	Purple	Shatter Resistance	Very Good	Phytophthora Field Tol.	Very Good				
Hilum Color	Imp.Black	Stress Tolerance	Excellent	SDS Tolerance	Very Good				
Pod Color	Tan	SCN Resistance Source	PI88788	White Mold	Very Good				
Pubescence	Gray	Phytophthora Res. Source	None	Stem Canker	Excellent				
Protein	NA	No-Till Adaptability	Excellent	Root Knot	NA				
Oil	NA			Cover Crop Companion	Fair				
Soil Placement		Recommended Seeding Rates				Herbicide Tolerance			
Light Soils: Low O.M., Low CEC, Low water holding capacity.	HR	Row Width	120/130	135/145	150/160	Conventional		Dicamba	
Medium Soils: O.M. 1.5-3.5%, CEC 11-20, Good productivity, well drained silt loams.	HR	30" Rows	R	HR	HR	Glyphosate		2,4-D	
Heavy Soils: High O.M. >3.5%, CEC >20, Well drained highly productive with deep top soil	HR	Twin Rows	HR	HR	R	Glufosinate		Callisto	
Poorly Drained Soils: Soils that tend to remain saturated for extended periods of time.	HR	20" Rows	R	R	HR	Sulfonurea		Isoxofluto	
		<20" rows	NR	R	HR				
<small>Population ranges are listed in 1000 seeds/acre</small>									
NR=Not Recommended, R=Recommended, HR=Highly Recommended									
Notes:									

